

- instability:comparison with magnetic resonance imaging [J].Injury, 2017,48(10):2323–2328.
- [12] Willekens I,Kichouh M,Boulet C,et al.Ultrasound follow-up of posttraumatic injuries of the sagittal band of the dorsal hood treated by a conservative approach[J].Eur J Radiol,2015,84(2):278–283.
- [13] Sato J,Ishii Y,Noguchi H.Diagnostic performance of the extensor carpi ulnaris (ECU) synergy test to detect sonographic ECU abnormalities in chronic dorsal ulnar-sided wrist pain [J].J Ultrasound Med,2016,35(1):7–14.
- [14] Ahmet S,Murat B,Duygu K.Intrauterine temporomandibular joint dislocation:prenatal sonographic evaluation[J].Clin Med Res,2014,12(1–2):58–60.
- [15] Trikha SP,Acton D,O'Reilly M,et al.Acute lateral dislocation of the patella:correlation of ultrasound scanning with operative findings[J].Injury,2003,34(8):568–571.
- [16] Shih JC,Peng SS,Hsiao SM,et al.Three-dimensional ultrasound diagnosis of Larsen syndrome with further characterization of neurological sequelae[J].Ultrasound Obstet Gynecol,2004,24(1):89–93.
- [17] Kolb A,Schweiger N,Mailath-Pokorny M,et al.Low incidence of early developmental dysplasia of the hip in universal ultrasonographic screening of newborns:analysis and evaluation of risk factors [J].Int Orthop,2016,40(1):123–127.
- [18] Tafazal S,Flowers MJ.Do we need to follow up an early normal ultrasound with a later plain radiograph in children with a family history of developmental dysplasia of the hip? [J].Eur J Orthop Surg Traumatol,2015,25(7):1171–1175.
- [19] 周小平,祝伟宏,石静.超声波在小婴儿髋关节脱位治疗中的监测作用[J].当代医学,2014,20(32):22–23.
- [20] Karnik AS,Karnik A,Joshi A.Ultrasound examination of pediatric musculoskeletal diseases and neonatal spine[J].Indian J Pediatr,2016,83(6):565–577.
- [21] Ömeroğlu H.Use of ultrasonography in developmental dysplasia of the hip[J].J Child Orthop,2014,8(2):105–113.
- [22] Upasani VV,Bomar JD,Matheney TH,et al.Evaluation of brace treatment for infant hip dislocation in a prospective cohort [J].J Bone Joint Surg Am,2016,98(14):1215–1221.
- [23] Fukuiage K,Futami T,Ogi Y,et al.Ultrasound-guided gradual reduction using flexion and abduction continuous traction for developmental dysplasia of the hip [J].Bone Joint J,2015,97(3):405–411.
- [24] Ömeroğlu H,Köse N,Akceylan A.Success of pavlik harness treatment decreases in patients  $\geq 4$  months and in ultrasonographically dislocated hips in developmental dysplasia of the hip [J].Clin Orthop Relat Res,2016,474(5):1146–1152.
- [25] David M,Robb C,Jawanda S,et al.Late recurrence of developmental dysplasia of the hip following Pavlik harness treatment until normal ultrasound appearance[J].J Orthop,2015,12(2):81–85.

(收稿日期:2017-03-15)

## · 病例报道 ·

## Ultrasonic manifestations of huge broad ligament with huge ovarian teratoma:a case report 巨大阔韧带肌瘤伴巨大卵巢畸胎瘤超声表现 1 例

曹小容

[中图法分类号]R445.1

[文献标识码] B

患者女,45岁,因“间断性尿频半年加重伴右侧腰部不适1d”就诊。超声检查:盆腹腔见一大小约19.0 cm×15.8 cm×9.7 cm巨大实质性包块,左侧卵巢大小形态正常,右侧卵巢未满意探及。右侧输尿管上段扩张伴右肾积水(图1,2);CDFI可探及包块内少许条状血流信号(图3)。超声提示:浆膜下子宫肌瘤可能性大。MRI检查:子宫右侧壁巨大肿块影,大小约23 cm×13 cm×8 cm,与子宫浆膜层分界不清,右后方见一大小约10 cm×5 cm×6 cm类圆形混杂信号。MRI提示:①子宫肌瘤(浆膜下)可能性大;

②考虑畸胎瘤。入院后2d行剖腹探查术,术中见右侧阔韧带内见一大小约25 cm×15 cm×10 cm巨大包块,其基底部与子宫峡部由一蒂相连,子宫被包块挤向左侧盆腔。右侧输尿管变形,沿包块外侧阔韧带基底部走行再向内而进入膀胱。右侧卵巢增大,约8 cm×5 cm×5 cm。行右侧输尿管镜下输尿管支架植入术、右侧阔韧带肌瘤摘除术、右侧附件切除术及子宫次全切除术,术后剖开包块内见灰白色水肿的肌性组织,螺旋状结构分界不清。

(下转123页)